

12. REFERENCES

NGĀ TOHUTORU

- Abookire, A. A., Duffy-Anderson, J. T., Jump, C. M. (2007). Habitat associations and diet of young-of-the-year Pacific cod (*Gadus macrocephalus*) near Kodiak, Alaska. *Marine Biology*, 150 (4): 713–726.
- Abraham, G. (2005). Holocene sediments of Tamaki Estuary: characterisation and impact of recent human activity on an urban estuary in Auckland, New Zealand. Unpublished PhD Thesis, University of Auckland, Auckland, New Zealand.
- Aguirre J.D., Bollard-Breen, B., Cameron, M., Constantine, R., Duffy, C.A.J., Dunphy, B., Hart, K., Hewitt, J.E., Jarvis, R.M., Jeffs, A., Kahui-McConnell, R., Kawharu, M., Liggins, L., Lohrer, A.M., Middleton, I., Oldman, J., Sewell, M.A., Smith, A.N.H., Thomas, D.B., Tuckey, B., Vaughan, M., Wilson, R. (in press). Loved to pieces: Toward the sustainable management of the Waitematā Harbour and Hauraki Gulf. *Regional Studies in Marine Science*. Available after 7 March 2016 from <http://dx.doi.org/10.1016/j.rsma.2016.02.009>.
- Airoldi, L., Beck, M.W. (2007). Loss, status and trends for coastal marine habitats of Europe. *Oceanography and Marine Biology: An Annual Review* 45: 345–405.
- Alfaro, A.C. (2006). Benthic macro-invertebrate community composition within a mangrove/seagrass estuary in northern New Zealand. *Estuarine, Coastal and Shelf Science*, 66(1–2): 97–110.
- Armstrong, C.W., Falk-Petersen, J. (2008). Habitat–fisheries interactions: a missing link? *ICES Journal of Marine Science*, 65: 817–821.
- Attrill, M.J., Strong, J.A., Rowden, A.A. (2000). Are macroinvertebrate communities influenced by seagrass structural complexity? *Ecography*, 23: 114–121.
- Auckland Council. (2007). Erosion and sediment control guide for land disturbing activities in the Auckland region. Technical Publication 90. Retrieved from <http://www.aucklandcouncil.govt.nz/EN/planspoliciesprojects/reports/technicalpublications/Pages/technicalpublications51-100.aspx>
- Auckland Council. (2014). On-site wastewater management. Retrieved from <http://www.aucklandcouncil.govt.nz/EN/ratesbuildingproperty/consents/buildingstructures/Documents/onsitewastewatermanagementintro.pdf>
- Auckland Council. (n.d.). Biodiversity on your property. Website at <http://www.aucklandcouncil.govt.nz/en/environmentwaste/biodiversity/pages/biodiversityonyourproperty.aspx>
- Auster, P.J., Langton, R.W. (1999). The effects of fishing on fish habitat. In Benaka, L.R. (Ed.), *Fish Habitat: Essential Fish Habitat and Rehabilitation*. American Fisheries Society Symposium 22. American Fisheries Society, Bethesda, Maryland. 150–187.
- Auster, P.J., Malatesta, R.J., Langton, R.W., Watling, L., Valentine, P.C., Donaldson C.L.S., Langton, E.W., Shepard, A.N., Babib, I.G. (1996). The impacts of mobile fishing gear on seafloor habitats in the Gulf of Maine (Northwest Atlantic): implications for conservation of fish populations. *Review in Fisheries Science*, 4: 185–202.
- Ayling, T., Schiel, D. (2003). Poor Knights Islands. In Andrew, N., & Francis, M. Craig (Eds.), *The Living Reef – the ecology of New Zealand’s rocky reefs*. Potton Publishing, Wellington.
- Babcock, R.C., Kelly, S., Shears, N.T., Walker, J.W., Willis, T.J. (1999). Changes in community structure in temperate marine reserves. *Marine Ecology Progress Series* 189: 125–134. Retrieved from www.int-res.com/articles/meps/189/m189p125.pdf
- Baillon, S., Hamel, J., Wareham, V., Mercier, A. (2012). Deep cold-water corals as nurseries for fish larvae. *Frontiers in Ecology and the Environment*, 10: 351–356.
- Baker, A.N., Madon, B. (2002). Bryde’s whales (*Balaenoptera cf. brydei* Olsen 1913) in the Hauraki Gulf and northeastern New Zealand waters. *Science for Conservation* 272. Department of Conservation, Wellington. Retrieved from <http://www.doc.org.nz/Documents/science-and-technical/sfc272.pdf>

- Baker, C. S., Chilvers, B.L., Constantine, R., DuFresne, S., Mattlin, R.H., van Helden, A., Hitchmough, R. (2010). Conservation status of New Zealand marine mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research*, 44: 101–115.
- Ball, B., Munday, B., Tuck, I. (2000). Effects of otter trawling on the benthos and environment in muddy sediments. In Kaiser, M.J., de Groot (Eds.), *Effects of Fishing on Non-target Species and Habitats*, pp. 69–82. S.J. Blackwell Science, Oxford. 399.
- Barbera, M. (2012). Towards an economic valuation of the Hauraki Gulf: A stock-take of activities and opportunities. Technical Report TR2012/035, Auckland Council, Auckland. 125.
- Battley, P.F., Brownell, B. (ed.) (2007). Population biology and foraging ecology of waders in the Firth of Thames - update 2007. Auckland Regional Council, Technical Publication No. 347.
- Beazley, L. I., Kenchington, E. L., Murillo, F. J., del Mar Sacau, M. (2013). Deep-sea sponge grounds enhance diversity and abundance of epibenthic megafauna in the Northwest Atlantic. *ICES Journal of Marine Science*, 70: 1471–1490.
- Beck, M.W., Heck, Jr. K.L., Able, K.W., Childers, D.L., Eggleston, D.B., Gillanders, B.M., Halpern, B., Hays, C.G., Hoshino, K., Minello, T.J., Orth, R.J., Sheridan, P.F., Weinstein, M.P. (2001). The identification, conservation, and management of estuarine and marine nurseries for fish and invertebrates. *BioScience*, 51(8): 633–641.
- Beentjes, M.P., Francis, M.P. (1999). Movement of hapuka (*Polyprion oxygeneios*) determined from tagging studies. *New Zealand Journal of Marine and Freshwater Research*, 33(1): 1–12.
- Bell, J.D., Galzin, R. (1984). Influence of live coral cover on coral-reef fish communities. *Marine Ecology Progress Series*, 15: 265–274.
- Bell, S.S., Hicks, G.R.F. (1991). Marine landscapes and faunal recruitment: a field test with seagrasses and copepods. *Marine Ecology Progress Series*, 73: 61–68.
- Bentley, N., Davies, N.M., McNeill, S.E. (2004). A spatially explicit model of the snapper (*Pagrus auratus*) fishery in SNA1. *New Zealand Fisheries Assessment Report 2004/26*.
- Berghan, J., Algie, K. D., Stockin, K. A., Wiseman, N., Constantine, R., Tezanos-Pinto, G., Mourão, F. (2008). A preliminary photo-identification study of bottlenose dolphin (*Tursiops truncatus*) in Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 42: 465–472.
- Beukers-Stewart, B.D., Vause, B.J., Mosley, M.W.J., Rossetti, H.L., & Brand, A.R. (2005). Benefits of closed area protection for a population of scallops. *Marine Ecology Progress Series* 298: 189–204.
- Booth, J.D. (1984). Movements of packhorse rock lobsters (*Jasus verreauxi*) tagged along the eastern coast of the North Island, New Zealand. *New Zealand Journal of Marine & Freshwater Research*, 18: 275–281.
- Booth, J.D. (1997). Long-distance movements in *Jasus* spp. and their role in larval recruitment. *Bulletin of Marine Science*, 61: 111–128.
- Boström, C., Jackson, E.L., Simenstad, C.A. (2006). Seagrass landscapes and their effects on associated fauna: a review. *Estuarine, Coastal and Shelf Science*, 68(3): 383–403.
- Bracken, M.E.S., Braken, B.E., Rogers-Bennett, L. (2007). Biodiversity, foundation species, and marine ecosystem management. *California Cooperative Oceanic Fisheries Investigations Report* 48.
- Bradford-Grieve, J., Probert, K., Lewis, K., Sutton, P., Zeldis, J., Orpin, A. (2006). Chapter 36. New Zealand shelf region. In Robinson, A, R., Brink, K. H. (Eds.), *The Sea: Ideas and Observations on Progress in the Study of Seas Vol. 14. Interdisciplinary Regional Studies and Syntheses. Part B*. Wiley, New York. 1451-1492
- Brasher, D.J., Ovenden, J.R., Booth, J.D., White, R.G.W. (1992). Genetic subdivision of Australian and New Zealand populations of *Jasus verreauxi* (Decapoda: Palinuridae)—preliminary evidence from the mitochondrial genome. *New Zealand Journal of Marine and Freshwater Research*, 26(1): 53–58.

- Buchsbaum, R., Pederson, J., Robinson, W.E. (2005). The decline of fisheries resources in New England: evaluating the impact of overfishing, contamination, and habitat degradation. MIT Sea Grant College Program Publication No. 05–5.
- Buhl-Mortensen, L., Vanreusel, A., Gooday, A.J., Levin, L.A., Priede, I.G., Buhl-Mortensen, P., Gheerardyn, H., King, N.J., Raes, M. (2010). Biological structures as a source of habitat heterogeneity and biodiversity on the deep ocean margins. *Marine Ecology*, 31: 21–50.
- Caddy, J. F., Defeo, O. (2003). Enhancing or restoring the productivity of natural populations of shellfish and other marine invertebrate resources. FAO Fisheries Technical Paper 448. Mexico.
- Carr, M. H. (1989). Effects of macroalgal assemblages on the recruitment of temperate zone reef fishes. *Journal of Experimental Marine Biology and Ecology*, 126: 59–76.
- Chang, F.H., Zeldis, J., Gall, M., Hall, J.A. (2003). Seasonal and spatial variation of phytoplankton functional groups on the northeastern New Zealand continental shelf and in Hauraki Gulf. *Journal of Plankton Research*, 25: 737–758.
- Charton, J.A.G., Ruzafa, A.P. (1998). Correlation between habitat structure and a rocky reef fish assemblage in the Southwest Mediterranean. *Marine Ecology*, 19: 111–128.
- Chiaroni, L., Hewitt, J.E., Hancock, N. (2008). Benthic marine habitats and communities of Kawau Bay. Prepared by NIWA for Auckland Regional Council. Auckland Regional Council Technical Report 2008/006.
- Claudia Hellberg, Ian Mayhew and David Mead 2016. Auckland’s stormwater management under the Auckland Unitary Plan. Proceedings, Stormwater Conference 2016, Water New Zealand, 18–20 May 2016, Nelson.
- Cloern, J.E. (2001). Our evolving conceptual model of the coastal eutrophication problem. *Marine Ecology Progress Series* 210: 223–253.
- Cole, R.G., Villouta, E., Davidson, R.J. (2000). Direct evidence of limited dispersal of the reef fish *Parapercis colias* (Pinguipedidae) within a marine reserve and adjacent fished areas. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 10(6): 421–436.
- Collie, J.S., Escanero, G.A., Valentine, P.C. (1997). Effects of bottom fishing on the benthic megafauna of Georges Bank. *Marine Ecology Progress Series* 155: 159–172.
- Collie, J.S., Escanero, G.A., Valentine, P.C. (2000a). Photographic evaluation of the impacts of bottom fishing on benthic epifauna. *ICES Journal of Marine Science*, 57: 987–1001.
- Collie, J.S., Hall, S.J., Kaiser, M.J., Poiner, I.R. (2000b). A quantitative analysis of fishing impacts on shelf-sea benthos. *Journal of Animal Ecology*, 69: 785–798.
- Connell, J.H. (1978). Diversity in tropical rain forests and coral reefs. *Science*, 199: 1302–1310.
- Connell, S.P., Jones, G.P. (1991). The influence of habitat complexity on post-recruitment processes in a temperate reef fish population. *Journal of Experimental Marine Biology and Ecology*, 151: 271–294.
- Connolly, R.M., Hindell, J.S., Gorman, D. (2005). Seagrass and epiphytic algae support the nutrition of a fisheries species, *Sillago schomburgkii*, in adjacent intertidal habitats. *Marine Ecology Progress Series* 286: 69–79.
- Constantine, R. (2002). The Behavioural Ecology of the Bottlenose Dolphins (*Tursiops truncatus*) of Northeastern New Zealand: A Population Exposed to Tourism. PhD. thesis, School of Biological Sciences, The University of Auckland.
- Coriolis Research. (2012). Industry investment opportunities in the New Zealand salmon industry. Retrieved from <http://www.mbie.govt.nz/info-services/sectors-industries/food-beverage/documents-image-library/Investment%20opportunities%20in%20the%20salmon%20industry%20-PDF%201.2%20MB.pdf>
- Costanza, R., d’Arge, R., de Groot, R., Farber, S., Grasso, M.B., Hannon, B., Limburg, K., Naeem, S., O’Neill, R.V. Paruelo, J. Raskin, R.G., Sutton, P., & Belt, v.d.M. (1997). The value of the world’s ecosystem services and natural capital. *Nature*, 387: 253–260.

- Craggs, R., Park, J., Heubeck, S. and Sutherland, D. (2014). High rate algal pond systems for low-energy wastewater treatment, nutrient recovery and energy production. *New Zealand Journal of Botany*, 52(1): 60–73.
- Craggs, R., Park, J., Sutherland, D. and Heubeck, S. (2015). Economic construction and operation of hectare-scale wastewater treatment enhanced pond systems. *Journal of Applied Phycology*, 27: 1913–1922.
- Cromarty, P. and Scott, D.A. (Eds.). (1995). *A Directory of Wetlands in New Zealand*. Department of Conservation, Wellington. Retrieved from <http://www.doc.govt.nz/Documents/science-and-technical/nzwetlands00.pdf>
- Crossland, J. (1981). *Fish Eggs and Larvae of the Hauraki Gulf, New Zealand*. Fisheries Research Bulletin No. 23. Fisheries Research Division, New Zealand Ministry of Agriculture and Fisheries, Wellington.
- Cummings, V.J., Thrush, S.F., Hewitt, J.E., Funnell, G.A. (2001). Variable effect of a large suspension-feeding bivalve on infauna: experimenting in a complex system. *Marine Ecology Progress Series* 209: 159–175.
- Dawbin, W.H. (1966). The seasonal migratory cycle of humpback whales. In: Norris KS (ed) *Whales, dolphins, and porpoises*. University of California Press, Berkeley and Los Angeles, 145–171.
- Dayton, P.K., Tegner, M.J., Edwards, P.B., Riser, K.L. (1998). Sliding baselines, ghosts, and reduced expectations in kelp forest communities. *Ecological Applications*, 8(2): 309–322.
- Dayton, P.K., Thrush, S., Coleman, F.C. (2002). The ecological effects of fishing in marine ecosystems of the United States. Pew Oceans Commission, Arlington, VA.
- Dayton, P.K., Thrush, S.F., Agardy, T.M., Hofman, R.J. (1995). Environmental effects of fishing. *Aquatic Conservation: Marine and Freshwater Ecosystems* 5: 205–232.
- Dean, R.L., Connell, J.H. (1987). Marine invertebrates in an algal succession. III. Mechanisms linking habitat complexity with diversity. *Journal of Experimental Marine Biology and Ecology*, 109: 249–273.
- Denham, R.N., Bannister, R.W., Guthrie, K.M., & Crook, F.G. (1984). Surveys of the East Auckland and East Cape currents, New Zealand. *Australian Journal of Marine and Freshwater Research*, 35: 491–504.
- Dennison, W.C., Orth, R.J., Moore, K.A., Stevenson, J.C., Carter, V., Kollar, S., Bergstrom, P.W., Batiuk, R.A. (1993). Assessing water quality with submersed aquatic vegetation. *BioScience*, 43: 86–94.
- Denny, C.M., Babcock, R.C. (2004). Do partial marine reserves protect reef fish assemblages? *Biological Conservation*, 116(1): 119–129.
- Department of Conservation. (n.d.). *Wetlands*. Retrieved from <http://www.doc.govt.nz/nature/habitats/wetlands/>
- Department of Fisheries and Oceans. (2006). Impacts of trawl gears and scallop dredges on benthic habitats, populations and communities. Canadian Science Advisory Secretariat Science Advisory Report 2006/025.
- Dromgoole, F., Foster, B. (1983). Changes to the marine biota of the Auckland Harbour. *Tane*, 29: 79–96.
- Ebeling, A.W., Laur, D.R. (1985). The influence of plant cover on surfperch abundance at an offshore temperate reef. *Environmental Biology of Fishes*, 12(3): 169–179.
- Eckman, J.E. (1987). The role of hydrodynamics in recruitment, growth, and survival of *Argopecten irradians* (L.) and *Anomia simplex* (D’Orbigny) within eelgrass meadows. *Journal of Experimental Marine Biology and Ecology*, 106: 165–191.
- Ellis J., Cummings V., Hewitt J., Thrush S., Norkko A. (2002). Determining effects of suspended sediment on condition of a suspension feeding bivalve (*Atrina zelandica*): Results of a survey, a laboratory experiment and a field transplant experiment. *Journal of Experimental Marine Biology and Ecology*, 267(2): 147–174.
- Ellis, J., Nicholls, P., Craggs, R., Hofstra, D., Hewitt, J. (2004). Effects of terrigenous sedimentation on mangrove physiology and associated macrobenthic communities. *Marine Ecology Progress Series*, 270: 71–82.

- Environment Foundation. (2015). Case Study: Auckland's Sustainable Catchments Programme. Auckland. Retrieved from <http://www.environmentguide.org.nz/issues/marine/catchment-based-activities/im:2133/>
- Fernandes, L, Green, A., Tanzer, J., White, A., Alino, P. M., Jompa, J., Lokani, P., Soemodinoto, A., Knight, M., Pomeroy, B., Possingham, H., Pressey, B. (2012). Biophysical principles for designing resilient networks of marine protected areas to integrate fisheries, biodiversity and climate change objectives in the Coral Triangle. Report prepared by The Nature Conservancy for the Coral Triangle Support Partnership. 152.
- Fonseca, M.S., Koehl, M.A.R. (2006). Flow in seagrass canopies: The influence of patch width. *Estuarine, Coastal and Shelf Science*, 67: 1–9.
- Francis, M.P. (2010). Movement of tagged rig and school shark among QMAs, and implications for stock management boundaries. New Zealand Fisheries Assessment Report 2010/3. 24. Ministry for Primary Industries, Wellington. Retrieved from http://fs.fish.govt.nz/Doc/22148/10_03_FAR.pdf.ashx
- Francis, M.P., Morrison, M.A., Leathwick, J., Walsh, C. (2011). Predicting patterns of richness, occurrence and abundance of small fish in New Zealand estuaries. *Marine and Freshwater Research*, 62: 1327–1341.
- Francis, M.P., Morrison, M.A., Leathwick, J., Walsh, C., Middleton, C. (2005). Predictive models of small fish presence and abundance in northern New Zealand harbours. *Estuarine, Coastal and Shelf Science*, 64: 419–435.
- Francis, M.P., Mulligan, K.P., Davies N.M., & Beentjes, M.P. (1999). Age and growth estimates for New Zealand hapuka, *Polyprion oxygeneios*. *Fishery Bulletin*, 97(2): 227–242.
- Gacia, E., Duarte, C.M. (2001). Sediment retention by a Mediterranean *Posidonia oceanica* meadow: the balance between deposition and resuspension. *Estuarine, Coastal and Shelf Science*, 52: 505–514.
- Garrigue, C., Zerbini, A. N., Geyer, Y., Heide-Jørgensen, M., Hanaoka, W., Clapham, P. (2010). Movements of satellite-monitored humpback whales from New Caledonia. *Journal of Mammalogy*, 91(1): 109–115.
- Gaskin, C., Rayner, M. (2013). Seabirds of the Hauraki Gulf: Natural History, Research and Conservation. Hauraki Gulf Forum. 143. Retrieved from <http://www.aucklandcouncil.govt.nz/EN/AboutCouncil/representativesbodies/haurakigulfforum/Documents/haurakigulfseabirdmanagementplanmar2013.pdf>
- Gilbert, C.S, Gentleman, W.C., Johnson, C.L., DiBacco, C., Pringle, J.M., & Chen, C. (2010). Modelling dispersal of sea scallop (*Placopecten magellanicus*) larvae on Georges Bank: The influence of depth-distribution, planktonic duration and spawning seasonality. *Progress in Oceanography*, 87(1–4): 37–48.
- Gislason, H. (1994). Ecosystem effects of fishing activities in the North Sea. *Marine Pollution Bulletin*, 29: 520–527.
- Gislason, H., Sinclair, M., Sainsbury, K., O'Boyle, R. (2000). Symposium overview: incorporating ecosystem objectives within fisheries management. *ICES Journal of Marine Science*, 55: 362–370.
- Graham, D.H. (1939). Breeding habits of the fishes of Otago Harbour and adjacent seas. *Transactions of the Royal Society of New Zealand*, 69: 361–372.
- Gratwike, B., Speight, M.R. (2005). The relationship between fish species richness, abundance and habitat complexity in a range of shallow tropical marine environments. *Journal of Fish Biology*, 66: 650–667.
- Grech, A., Chartrand-Miller, K., Erftemeijer, P., Fonseca, M., McKenzie, L., Rasheed, M., Coles, R. (2012). A comparison of threats, vulnerabilities and management approaches in global seagrass bioregions. *Environmental Research Letters*, 7(2): 4006.
- Green, E.P., Short, F.T. (Eds.), (2003). *World atlas of seagrasses*. University of California Press, Berkeley, California. 324.
- Green, M.O., Zeldis, J. (2015). Firth of Thames Water Quality and Ecosystem Health – A Synthesis. NIWA Client Report HAM2015–016, April 2015. 81. Retrieved from <http://www.waikatoregion.govt.nz/tr201523/>
- Greenway, J.P.C. (1969). Survey of mussels (Mollusca: Lamellibranchia) in the Firth of Thames, 1961–67. *New Zealand Journal of Marine & Freshwater Research*, 3: 304–317.

- Hadfield, M., O'Callaghan, J., Pritchard, M., Stevens, C. (2014). Sediment transport and deposition in the Hauraki Gulf: a pilot modelling study. NIWA Client Report No: WLG2012-29. National Institute of Water & Atmospheric Research Ltd., Wellington. 28.
- Hall, J.A., Safi, K., James, M.R., Zeldis, J., Weatherhead, M. (2006). Microbial assemblage dynamics during the spring-summer transition on the northeast continental shelf of New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 40: 195–210.
- Hall, S.J. (1999). The effects of fisheries on ecosystems and communities (Fish and Aquatic Resources). Blackwell Scientific, Oxford, England. 274.
- Hall, S.J. (1999). The effects of fishing on marine ecosystems and communities (Fish Biology and Aquatic Resources Series 1). Blackwell Science, Oxford. 274.
- Hartill, B.W., Payne, G.W., Rush, N., & Bian, R. (2016). Bridging the temporal gap: Continuous and cost-effective monitoring of dynamic recreational fisheries by web cameras and creel surveys. *Fisheries Research*, 183: 488–497
- Hauraki Gulf Forum. (2011). State of our Gulf: Tīkapa Moana Hauraki Gulf state of the environment report 2011. Hauraki Gulf Forum, Auckland. 163.
- Hay, R. (1983). Shorebirds of the Miranda coastline. *Tane*, 29: 15–30.
- Hayward, B.W., Stephenson, A.B., Morley, M., Riley, J.L., Grenfell, H.R. (1997). Faunal changes in Waitematā Harbour sediments, 1930s–1990s. *Journal of the Royal Society of New Zealand*, 27(1): 1–20.
- Heck Jr, K.L., Hays, G., Orth, R.J. (2003). Critical evaluation of the nursery role hypothesis for seagrass meadows. *Marine Ecology Progress Series*, 253: 123–136.
- Heck, K.L., Wetstone, G.S. (1977). Habitat complexity and invertebrate species richness and abundance in tropical seagrass meadows. *Journal of Biogeography*, 4: 135–142.
- Heifetz, J. (2002). Coral in Alaska: distribution, abundance, and species associations. *Hydrobiologia*, 471: 19–28.
- Hemminga, M.A., Duarte, C.M. (2000). Elemental dynamics in seagrass systems. In M.A. Hemminga and C.M. Duarte (Eds.), *Seagrass Ecology*. Cambridge University Press. 146–189.
- Hemminga, M.A., Duarte, C.M. (Eds.), (2000). *Seagrass Ecology*, Cambridge University Press. 298.
- Henriques, P.R. (1980). Faunal community structure of eight soft shore, intertidal habitats in the Manukau Harbour. *New Zealand Journal of Ecology*, 3: 97–103.
- Hewitt J.E., Pilditch C.A. (2004). Environmental history and physiological state influence feeding responses of *Atrina zelandica* to suspended sediment concentrations. *Journal of Experimental Marine Biology and Ecology*, 306(1): 95–112.
- Hewitt, J.E., Thrush, S.F., Halliday, J., Duffy, C. (2005). The importance of small-scale habitat structure for maintaining beta diversity. *Ecology*, 86: 1619–1626.
- Hicks, G.R.F. (1986). Distribution and behaviour of meiofaunal copepods inside and outside seagrass beds. *Marine Ecology Progress Series*, 31: 159–170.
- Hicks, G.R.F. (1989). Does epibenthic structure negatively affect meiofauna? *Journal of Experimental Marine Biology and Ecology*, 133: 39–55.
- Hiddink, J.G., Jennings, S., Kaiser, M.J., Queiros, A.M., Duplisea, D.E., Piet, G.J. (2006). Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal Fisheries and Aquatic Science*, 63: 721–736.
- Hindell, J.S., Jenkins, G.P., Keough, M.J. (2000). Evaluating the impact of predation by fish on the assemblage structure of fishes associated with seagrass (*Heterozostera tasmanica*) (*Martens ex Ascherson*) den Hartog, and unvegetated sand habitats. *Journal of Experimental Marine Biology and Ecology*, 255: 153–174.

- Hindell, J.S., Jenkins, G.P., Keough, M.J. (2001). Spatial and temporal variability in the effects of fish predation on macrofauna in relation to habitat complexity and cage effects. *Marine Ecology Progress Series*, 224: 231–250.
- Hixon, M. A., Johnson, D.W., & Sogard, S. M. (2014). BOFFFFs: on the importance of conserving old-growth age structure in fishery populations. *ICES Journal of Marine Science*, 71: 2171–2185.
- Hobday, A.J., Tegner, M.J., & Haaker, P.L. (2000). Over-exploitation of a broadcast spawning marine invertebrate: Decline of the white abalone. *Reviews in Fish Biology and Fisheries*, 10: 493–514.
- Horinouchi, M. (2007). Review of the effects of within-patch scale structural complexity on seagrass fishes. *Journal of Experimental Marine Biology and Ecology*, 350(1): 111–129.
- Hume, T.M., McGlone, M.S. (1986). Sedimentation patterns and catchment use change recorded in the sediments of a shallow tidal creek, Lucas Creek, upper Waitematā Harbour, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 20: 677–687.
- Humphries, A. T., La Peyre, M.K., Kimball, M.E., Rozas, L.P. (2011). Testing the effect of habitat structure and complexity on nekton assemblages using experimental oyster reefs. *Journal of Experimental Marine Biology and Ecology*, 409(1–2): 172–179.
- Inglis, G. (2003). The Seagrasses of New Zealand. In Green, E.P., Short, F.T. (Eds.), *World Atlas of Seagrasses*. 134–143. University of California Press.
- IUCN World Commission on Protected Areas (IUCN-WCPA). (2008). *Establishing Marine Protected Area Networks—Making It Happen*. Washington, D.C.: IUCN-WCPA, National Oceanic and Atmospheric Administration and The Nature Conservancy. 118.
- Ivanovic, A., Zhu, J., Neilson, R., O'Neill, F. (2011). Modelling the physical impact of trawl components on the seabed and comparison with sea trials. *Ocean Engineering*, 38: 925–933.
- Jackson, J.B.C. (2001). What was natural in the coastal oceans? *Proceedings of the Natural Academy of Sciences of the United States of America*, 98(10): 5411–5418.
- Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R.H., Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, K.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., Warner, R.R. (2001). Historical overfishing and the recent collapse of coastal ecosystems. *Science*, 293(5530): 629–637.
- Jackson, S. E. (2014). *Prioritisation of Areas in the Hauraki Gulf Marine Park for Biodiversity Conservation*. Masters of Science thesis in Marine Science, The University of Auckland. 112.
- Jennings, S., Dinmore, T.A., Duplisea, D.E., Warr, K.J., Lancaster, J.E. (2001). Trawling disturbance can modify benthic production processes. *Journal of Animal Ecology*, 70: 459–475.
- Jennings, S., Kaiser, M.J. (1998). The effects of fishing on marine ecosystems. *Advances in Marine Biology*, 34: 203–314.
- Johnson, P. and Gerbeaux, P. (2004). *Wetland Types in New Zealand*. Department of Conservation, Wellington.
- Jones, E.G., Morrison, M.A., Davey, N., Mills, S., Pallentin, A., Shankar, U., George, S., Kelly, M., Shankar, U. (In review). Biogenic habitats on the continental shelf. Part II: National field survey and analysis. *New Zealand Aquatic Environment and Biodiversity Report*. Wellington: Ministry for Primary Industries.
- Jones, E.G., Morrison, M.A., Davey, N., Hartill, B.W., Sutton, C. (in press). Biogenic habitats on New Zealand's continental shelf. Part I: Local Ecological Knowledge. *New Zealand Aquatic Environment and Biodiversity Report*. Ministry for Primary Industries, Wellington.
- Jones, E.G., Parsons, D., Morrison, M., Bagley, N., Paterson, C., Usmar, N. (2010). Chapter 13: Fish communities. In: *Bay of Islands OS20/20 survey report, NIWA Client Report: WLG2010–38*.

- Kahui - McConnell, R. (2012). Ōkahu Catchment Ecological Restoration Plan. Auckland: Ngāti Whātua Ōrākei
- Kaiser, M.J. (1998). Significance of bottom-fishing disturbance. *Conservation Biology*, 12: 1230–1235.
- Kaiser, M.J., Clarke, K.R., Hinz, H., Austen, M.C., Somerfield, P.J., Karakassis, I. (2006). Global analysis of response and recovery of benthic biota to fishing. *Marine Ecology Progress Series*, 311: 1–14.
- Kaiser, M.J., de Groot, S.J. (Eds.). (2000). *The effects of fishing on non-target species and habitats*. Blackwell Science, Oxford, England.
- Kelly, S., Sim-Smith, C., Faire, S., Pierre, J., & Hikuroa, D. C. H. (2014). State of our Gulf 2014, Hauraki Gulf - Tīkapa Moana / Te Moananui-ā-Toi, State of the Environment Report 2014 (4). Auckland Council: Hauraki Gulf Forum. Retrieved from <http://www.aucklandcouncil.govt.nz/EN/environmentwaste/coastalmarine/Documents/stateofourgulf2014.pdf>
- Kensler, C.B. (1967). The distribution of spiny lobsters (Crustacea: Decapoda: Palinuridae) in New Zealand waters. *New Zealand Journal of Marine & Freshwater Research*, 1: 412–420.
- Keough, M.J., Jenkins, G.P. (1995). Seagrass meadows and their inhabitants. In A.J., Chapman, M.G. Chapman (Eds.), *Underwood, Coastal marine ecology of temperate Australia*. UNSW Press, Sydney, Australia. 221–239.
- Klitgaard, A. (1995). The fauna associated with outer shelf and upper slope sponges (Porifera, Demospongiae) at the Faroe Islands, North-eastern Atlantic. *Sarsia*, 80: 1–22.
- Laman, E.A., Kotwicki, S., Rooper, C.N. (2015). Correlating environmental and biogenic factors with abundance and distribution of Pacific ocean perch (*Sebastes alutus*) in the Aleutian Islands. *Alaska Fisheries Bulletin*, 113: 270–289.
- Langlois, T.J., Anderson, M.J., Babcock, R.C. (2005). Reef-associated predators influence adjacent soft-sediment communities. *Ecology*, 86: 1508–1519.
- Larcombe, M.F. (1973). *Ecological report on the Waitematā Harbour*. Auckland Regional Authority, Auckland. 375.
- Leathwick, J.R., Rowden, A., Nodder, S., Gorman, R., Bardsley, S., Pinkerton, M., Baird, S.J., Hadfield, M., Currie, K., Goh, A. (2012). *A Benthic-optimised Marine Environment Classification (BOMECE) for New Zealand waters*. New Zealand Aquatic Environment and Biodiversity Report No. 88. Wellington: Ministry for Primary Industries. 54.
- Lee, S.T., Kelly, M., Langlois, T.J., Costello, M.J. (2015). Baseline seabed habitat and biotope mapping for a proposed marine reserve. *PeerJ*, 3:e1446 Retrieved from <https://doi.org/10.7717/peerj.1446>
- Leleu, K., Remy-Zephir, B., Grace, R., Costello, M.J. (2012). Mapping habitats in a marine reserve showed how a 30-year trophic cascade altered ecosystem structure. *Biological Conservation*, 155: 193–201.
- LePort, A., Montgomery, J.C., Croucher, A.E. (2014). Biophysical modelling of snapper *Pagrus auratus* larval dispersal from a temperate MPA. *Marine Ecology Progress Series*, 515: 203–215.
- Lindeboom, H.J., de Groot, S.J. (Eds.), (1998). *Impact-II: The effect of different types of fisheries on the North Sea and Irish Sea benthic ecosystems*. Netherlands Institute for Sea Research, Texel, The Netherlands. 404.
- Lindholm, J.B., Auster, P.J., Kaufman, L.S. (1999). Habitat-mediated survivorship of juvenile (0-year) Atlantic cod *Gadus morhua*. *Marine Ecology Progress Series*, 180: 247–255.
- Lomovasky, B.J., Gamero, P.A., Romero, L., Firstater, F.N., Gammara Salazar, A., Hidalgo, F., Tarazona, J., & Iribarne, O.O. (2015). The role of *Argopecten purpuratus* shells structuring the soft bottom community in shallow waters of southern Peru. *Journal of Sea Research*, 106: 14–26.
- Lotze, H.K., Coll, M., Magera, A.M., Ward-Paige, C., Airoldi, L. (2011). Recovery of marine animal populations and ecosystems. *Trends in Ecology & Evolution*, 26(11): 595–605.
- Lotze, H.K., Lenihan, H.S., Bourque, B.J., Bradbury, R.H., Cooke, R.G., Kay, M.C., Kidwell, S.M., Kirby, M.X., Petersen, C.H., Jackson, J.B.C. (2006). Depletion, degradation, and recovery potential of estuaries and coastal seas. *Science*, 312

(5781): 1806–1809.

Lowe, M., Morrison, M. (2012). Waterview Connection Causeway Project. Baseline Marine Ecological Monitoring: Open mudflat/sandflat fish surveys. Report prepared for New Zealand Transport Agency. NIWA Client Report No: NTA12101. National Institute of Water & Atmospheric Research Ltd., Auckland. 67.

Lowe, M.L., Morrison, M.A, Taylor, R.B. (2015). Harmful effects of sediment-induced turbidity on juvenile fish in estuaries. *Marine Ecology Progress Series*, 539: 241–254.

Luckhurst, B.E., Luckhurst, K. (1978). Analysis of the influence of substrate variables on coral reef fish communities. *Marine Biology*, 49: 317–323.

MacDiarmid, A.B. (1989). Size at onset of maturity and size dependent reproductive output of female and male spiny lobsters *Jasus edwardsii* (Hutton) (Decapoda, Palinuridae) in northern New Zealand. *Journal of Experimental Marine Biology and Ecology*, 127: 229–243.

MacDiarmid, A.B., Abraham, E., Baker, C.S., Carroll, E., Chagué-Goff, C., Cleaver, P., Francis, M.P., Goff, J., Horn, P., Jackson, J.A., Lalas, C., Lorrey, A., Marriot, P., Maxwell, K., McFadgen, B., McKenzie, A., Neil, H., Parsons, D., Patenaude, N., Paton, D., Paul, L.J., Pitcher, T., Pinkerton, M.H., Smith, I., Smith, T.D., Stirling B. (2016). Taking Stock – the changes to New Zealand marine ecosystems since first human settlement: synthesis of major findings, and policy and management implications. New Zealand Aquatic Environment and Biodiversity Report No. 170. Wellington: Ministry for Primary Industries. 48.

MacDiarmid, A.B., Butler, M.J. (1999). Sperm economy and limitation in spiny lobsters. *Behavioural Ecology and Sociobiology*, 46: 14–24.

Marine Parks Authority. (2008). A review of benefits of marine protected areas and related zoning considerations. Marine Parks Authority New South Wales. 14.

Martell, S.J.D., Walters, C.J., Wallace, S.S. (2000). The use of Marine Protected Areas for conservation of lingcod (*Ophiodon elongatus*). *Bulletin of Marine Science*, 66(3): 729–743.

Matheson, F., Wadhwa, S. (2012). Seagrass in Porirua Harbour: preliminary assessment of restoration potential. NIWA report for Greater Wellington Regional Council. (Unpublished report, available from Greater Wellington Regional Council). 35.

Matheson, F.E., Reed, J., Dos Santos, V.M., Cummings, V., Mackay, G. (in prep.) Comparison of transplant methods for seagrass rehabilitation in a temperate estuary. *Restoration Ecology*.

Maxwell, K., MacDiarmid, A.B. (2016). Taking Stock: Oral histories about changes in marine fauna observed locally in the Hauraki Gulf and along the Otago-Catlins coast, New Zealand, 1926–2008. New Zealand Aquatic Environment and Biodiversity Report No. 173. Wellington: Ministry for Primary Industries. 45.

McCormick, M.I. (1989). Spatio-temporal patterns in the abundance and population structure of a large temperate reef fish. *Marine Ecology Progress Series*, 53: 215–225.

McLeod, I.M., Parsons, D.M., Morrison, M., Le Port, A., Taylor, R.B. (2011). Factors affecting the recovery of soft-sediment mussel reefs in the Firth of Thames, New Zealand. *Marine and Freshwater Research*, 63: 78–83.

McLeod, I.M., Parsons, D.M., Morrison, M.A., Van Dijken, S.G., Taylor, R.B. (2014). Mussel reefs on soft sediments: a severely reduced but important habitat for macroinvertebrates and fishes in New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 48: 48–59.

Mendo, T., Moltschanivskyj, N., & Lyle, J.M. (2014). Role of density in aggregation patterns and synchronization of spawning in the hermaphroditic scallop *Pecten fumatus*. *Marine Biology*, 161: 2857–2868.

Millennium Ecosystem Assessment. (2005). Ecosystems and human well-being: biodiversity synthesis. World Resources

- Institute, Washington, D.C., USA. Retrieved from <http://www.unep.org/maweb/en/index.aspx>
- Mills, V.S., Berkenbusch, K. (2009). Seagrass (*Zostera muelleri*) patch size and spatial location influence infaunal macroinvertebrate assemblages. *Estuarine, Coastal and Shelf Science*, 81: 123–129.
- Ministry for Primary Industries. (2013). Review of sustainability measures and other management controls for SNA 1 for the 2013–14 fishing year. MPI Information Paper 2013/08, Ministry for Primary Industries, Wellington.
- Ministry for the Environment. (2003). Microbiological Water Quality Guidelines for Marine and Freshwater Recreational Areas. Wellington: MfE. Retrieved from <http://www.mfe.govt.nz/publications/fresh-water/microbiological-water-quality-guidelines-marine-and-freshwater-recreatio-13#twod4>
- Moffitt, E.A., Botsford, L.W., Kaplan, D.M., O'Farrell, M.R. (2009). Marine reserve networks for species that move within a home range. *Ecological Applications*, 19(7): 1835–1847.
- Morley, M. S., Hayward, B.W. (2009). Marine mollusca of Great Barrier Island, New Zealand. *Records of the Auckland Museum*, 46: 15–51.
- Morrison, M.A. (1990). Population dynamics of the scallop *Pecten novaezelandiae* in the Hauraki Gulf. Unpublished PhD thesis. University of Auckland.
- Morrison, M.A., Tuck, I.D., Taylor, R.B., Miller, A. (2016). An assessment of the Hauraki Gulf Cableway Protection Zone (CPZ), relative to adjacent seafloor. Auckland Council Technical Report: 54.
- Morrison, M.A., Jones, E., Consalvey, M., Berkenbusch, K. (2014). Linking marine fisheries species to biogenic habitats in New Zealand: a review and synthesis of knowledge. New Zealand Aquatic Environment and Biodiversity Report No. 130. Ministry for Primary Industries, Wellington. 156.
- Morrison, M.A., Jones, E.G., Parsons, D.P., Grant, C.M. (2014). Habitats and areas of particular significance for coastal finfish fisheries management in New Zealand: A review of concepts and life history knowledge, and suggestions for future research. New Zealand Aquatic Environment and Biodiversity Report No. 125. Ministry for Primary Industries, Wellington. 202.
- Morrison, M.A., Lowe, M.L., Grant, C.G., Smith, P.J., Carbines, G., Reed, J., Bury, S.J., Brown, J. (2014d). Seagrass meadows as biodiversity and productivity hotspots. New Zealand Aquatic Environment and Biodiversity Report No. 137. 147.
- Morrison, M.A., Lowe, M.L., Jones, E.G., Makey, L., Shankar, U., Usmar, N., Miller, A., Smith, M., Middleton, C. (2014c). Habitats of particular significance for fisheries management: The Kaipara Harbour. New Zealand Aquatic Environment and Biodiversity Report No. 129. Wellington: Ministry for Primary Industries. 169.
- Morrison, M.A., Lowe, M.L., Parsons, D.M., Usmar, N.R., McLeod, I.M. (2009). A review of land-based effects on coastal fisheries and supporting biodiversity in New Zealand. New Zealand Aquatic Environment and Biodiversity Report No. 37. Wellington: Ministry for Primary Industries. 100.
- Murray, C., and McDonald, G. (2010). Aquaculture: Economic impact in the Auckland region. Jointly prepared by the Auckland Regional Council and Market Economics Ltd for Auckland Regional Council.. Technical Report no. 009. Auckland Regional Council. Retrieved from <http://www.aucklandcouncil.govt.nz/SiteCollectionDocuments/aboutcouncil/planspublications/technicalpublications/tr2010009aquacultureeconomicimpactintheaucklandregion.pdf>
- National Institute for Water and Atmospheric Research. (2014). NIWA advances wastewater treatment processes. In *Freshwater and Estuaries Update*. NIWA. Retrieved from <https://www.niwa.co.nz/freshwater-and-estuaries/freshwater-and-estuaries-update/freshwater-update-62-september-2014/niwa-advances-wastewater-treatment>
- Nicholls, P., Hewitt, J., Halliday, J. (2003). Effects of suspended sediment concentrations on suspension and deposit feeding marine macrofauna. Auckland Regional Council Technical Publication 211. 40.
- Norkko, A., Hewitt, J.E., Thrush, S.F., Funnell, G.A. (2001). Benthic-pelagic coupling and suspension feeding bivalves:

- linking site-specific sediment flux and biodeposition to benthic community structure. *Limnology and Oceanography*, 46: 2067–2072.
- O’Callaghan, T.M., Baker, C.S. (2002). Summer cetacean community, with particular reference to Bryde’s whales, in the Hauraki Gulf, New Zealand. DOC Science Internal Series 55. Department of Conservation, Wellington. 18.
- Oldman, J., Hong, J., Stephens, S., Broekhuizen, N. (2007). Verification of Firth of Thames hydrodynamic model. Auckland Regional Council Technical Publication No. 326. 87.
- Oliver, W.R.B. (1923). Marine littoral plant and animal communities in New Zealand. *Transactions of the New Zealand Institute*, 54: 496–545.
- O’Neill, F.G., Summerbell, K. (2011). The mobilisation of sediment by demersal otter trawls. *Marine Pollution Bulletin*, 62(5): 1088-1097.
- Orth, R.J., Batiuk, R.A., Bergstrom, P.W., Moore, K.A. (2002). A perspective on two decades of policies and regulations influencing the protection and restoration of submerged aquatic vegetation in Chesapeake Bay, USA. *Bulletin of Marine Science*, 71: 1391–1403.
- Orth, R.J., Carruthers T.J.B., Dennison, W.B., Duarte, C.M., Fourqurean, J.W., Heck Jr., K.L., Hughes, A.R., Kendrick, G.A., Kenworthy, W.J., Olyarnik, S., Short, F.T., Waycott, M., Williams, S.L. (2006). A global crisis for seagrass ecosystems. *BioScience*, 56(12): 987–996.
- Palumbi, S.R., McLeod, K., Grumbaugh, D. (2008). Ecosystems in action: lessons from marine ecology about recovery, resistance, and reversibility. *BioScience*, 58: 33–42.
- Parsons, D., Morrison, M., Thrush, S., Middleton, C., Smith M., Spong K., Buckthought, D. (2013). The influence of habitat structure on juvenile fish in a New Zealand estuary. *Marine Ecology*, 34: 492–500.
- Parsons, D.M., Morrison, M.A., Slater, M.J. (2010). Responses to marine reserves: decreased dispersion of the sparid *Pagrus auratus* (snapper). *Biological Conservation*, 143: 2039–2048.
- Paul, L. (2012). A history of the Firth of Thames dredge fishery for mussels: use and abuse of a coastal resource. New Zealand Aquatic Environment And Biodiversity Report No. 94. Ministry for Primary Industries, Wellington.
- Paul, L.J. (2000). *New Zealand Fishes. Identification, natural history and fisheries*. Reed Books, Auckland, NZ. 253.
- Paul, L.J. (2002a). A description of the New Zealand fisheries for the two groper species, hapuka (*Polyprion oxygeneios*) and bass (*P. americanus*). New Zealand Fisheries Assessment Report 2002/13. Ministry for Primary Industries, Wellington. 47.
- Paul, L.J. (2002b). Can existing data describe the stock structure of the two New Zealand groper species, hapuka (*Polyprion oxygeneios*) and bass (*P. americanus*)? New Zealand Fisheries Assessment Report 2002/14. Ministry for Primary Industries, Wellington. 24.
- Pelletier, D., Mahévas, S. (2005). Fisheries simulation models for evaluating the impact of management policies, with emphasis on marine protected areas. *Fish and Fisheries*, 6(4): 307–349.
- Pérez-Matus, A., Shima, J. (2010). Disentangling the effects of macroalgae on the abundance of temperate reef fishes. *Journal of Experimental Marine Biology and Ecology* 388: 1-10.
- Peters, M., Clarkson, B. (Eds.), (2013). *Wetland Restoration: A Handbook for New Zealand Freshwater Systems*. Landcare and Research. Retrieved from <http://www.landcareresearch.co.nz/publications/books/wetlands-handbook>.
- Phillips, N.E., Shima, J.S. (2006). Differential effects of suspended sediments on larval survival and settlement of New Zealand urchins *Evechinus chloroticus* and abalone *Haliotis iris*. *Marine Ecology Progress Series*, 314: 149–158.
- Powell, A.W.B. (1937). Animal communities of the sea-bottom in Auckland and Manukau Harbours. *Royal Society of New Zealand Transactions and Proceedings*, 66: 354–401.

- Puig, P., Canals, M., Company, J.B., Martín, J., Amblas, D., Lastras, G., Palanques, A., Calafat, A. (2012). Ploughing the deep sea floor. *Nature*, 489: 286–289.
- Pusceddu, A., Bianchelli, S., Martín, J., Puig, P., Palanques, A., Masqué, P., Danovaro, R. (2014). Chronic and intensive bottom trawling impairs deep-sea biodiversity and ecosystem functioning. *Proceedings of the National Academy of Sciences*, 111: 8861–8866.
- Rabaut, M., Van de Moortel, L., Vincx, M., Degraer, S. (2010). Biogenic reefs as structuring factor in *Pleuronectes platessa* (Plaice) nursery. *Journal of Sea Research*, 64: 102–106.
- Radford, C.A., Sim-Smith, C.J., Jeffs, A.G. (2012). Can larval snapper, *Pagrus auratus*, smell their new home? *Marine and Freshwater Research*, 63: 898–904.
- Reed, J., Schwarz, A., Gosai, A., Morrison, M. (2004). Feasibility study to investigate the replenishment/reinstatement of seagrass beds in Whangarei Harbour. Phase 1. NIWA report for Northland Regional Council.
- Reid, B. (1969). Mussel survey Hauraki Gulf and Firth of Thames 1968. New Zealand Marine Department Fisheries Technical Report No. 34. 24.
- Rice, J.C. (2006). Impacts of Mobile Bottom Gears on Seafloor Habitats, Species, and Communities: A Review and Synthesis of Selected International Reviews. Canadian Science Advisory Secretariat Science Research Document 2006/057. 35.
- Roberts, C.M., Ormond, R.F.G. (1987). Habitat complexity and coral reef fish diversity and abundance on Red Sea fringing reefs. *Marine Ecology Progress Series*, 41: 1–8.
- Robertson, H.A., Dowding, J.E., Elliott, G.P., Hitchmough, R.A., Miskelly, C.M., O'Donnell, C.F.J., Powlesland, R.G., Sagar, P.M., Scofield, R.P., Taylor, G.A. (2013). Conservation status of New Zealand birds, 2012. New Zealand Threat Classification Series 4. New Zealand Department of Conservation. 22.
- Roemmich, D., Sutton, P. (1998). The mean and variability of ocean circulation past northern New Zealand: determining the representativeness of hydrographic climatologies. *Journal of Geophysical Research*, 103: 13041–13054.
- Rogers, A., Blanchard, J.L., Mumby, P.J. (2014). Vulnerability of coral reef fisheries to a loss of structural complexity. *Current Biology*, 24: 1000–1005.
- Rooker, J.R., Holt, G.J., Holt, S.A. (1998). Vulnerability of newly settled red drum (*Scianops ocellatus*) to predatory fish: Is early-life survival enhanced by seagrass meadows? *Marine Biology*, 131: 145–151.
- Sáenz-Arroyo, A., Roberts, C.M., Torre, J., Cariño-Olvera, M. (2005a). Using fishers' anecdotes, naturalists' observations and grey literature to reassess marine species at risk: the case of the Gulf grouper in the Gulf of California, Mexico. *Fish and Fisheries*, 6: 121–133.
- Sainsbury, K.J. (1987). Assessment and management of the demersal fishery on the continental shelf of north-western Australia. In Polovina, J.J., Ralston, S. (Eds.), *Tropical snappers and groupers: biology and fisheries management*. Westview Press, Boulder, CO. 465–503.
- Sainsbury, K.J. (1988). The ecological basis of multispecies fisheries and management of a demersal fishery in tropical Australia, In *Fish population dynamics* (2nd ed.). Gulland, I.A. (Eds.), John Wiley and Sons, Chichester and New York. 349–382.
- Sainsbury, K.J. (1991). Application of an experimental approach to management of a tropical multispecies fishery with highly uncertain dynamics. *ICES Marine Science Symposium*, 193: 301–320.
- Sainsbury, K.J., Campbell, R.A., Lindholm, R., Whitelaw, A.W. (1988). Experimental management of an Australian multispecies fishery: examining the possibility of trawl-induced habitat modification. *Global Trends Fisheries Management*, 20: 107–122.

- Sala, E., Knowlton, K. (2006). Global marine biodiversity trends. *Annual Review of Environment and Resources*, 31: 93–122.
- Salomon, A.K., Shears, N.T., Langlois, T.J., Babcock, R.C. (2008). Cascading effects of fishing can alter carbon flow through a temperate coastal ecosystem. *Ecological Applications*, 18(8): 1874–1887.
- Schwarz, A., Morrison, M., Hawes, I., Halliday, J. (2006). Physical and biological characteristics of a rare marine habitat: sub-tidal seagrass beds of offshore islands. Department of Conservation, Wellington. Science for Conservation No. 269. 39.
- Sharples, J., Greig, M.J.N. (1998). Tidal currents, mean flows and upwelling on the northeast shelf of New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 32(2): 215–231.
- Shears, N.T. (2010). Meola Reef ecological monitoring program: 2001–2010. Prepared by Auckland UniServices for Auckland Regional Council. Auckland Regional Council Technical Report 2010/031.
- Shears, N.T., Babcock, R.C. (2002). Marine reserves demonstrate top-down control of community structure on temperate reefs. *Oecologia*, 132: 131–142.
- Shears, N.T., Babcock, R.C. (2003). Continuing trophic cascade effects after 25 years of no take marine reserve protection. *Marine Ecology Progressive Series*, 246: 1–16.
- Shears, N.T., Babcock, R.C. (2007). Quantitative description of mainland New Zealand's shallow subtidal reef communities. Science for Conservation 280. Department of Conservation, Wellington. 126.
- Shears, N.T., Babcock, R.C., Salomon, A.K. (2008). Context-dependent effects of fishing: variation in trophic cascades across environmental gradients. *Ecological Applications*, 18(8): 1860–1873.
- Shears, N.T., Smith, F., Babcock, R., Duffy, C.A.J., Villouta, E. (2008). Evaluation of biogeographic classification schemes for conservation planning: application to New Zealand's coastal marine environment. *Conservation Biology*, 22(2): 467–481.
- Sheffield, A.T., Healy, T.R., McGlone, M.S. (1995). Infilling rates of a steep land catchment estuary, Whangamata, New Zealand. *Journal of Coastal Research*, 11(4): 1294–1308.
- Short, F.T., Polidoro, B., Livingstone, S.R., Carpenter, K.E., Bandeira, S., Bujang, J.S., Zieman, J.C. (2011). Extinction risk assessment of the world's seagrass species. *Biological Conservation*, 144(7): 1961–1971.
- Sim-Smith, C.J., Jeffs, A.G., Radford, C.A. (2012). Variation in the growth of larval and juvenile snapper, *Chrysophrys auratus* (Sparidae). *Marine & Freshwater Research*, 63: 1231–1243.
- Sim-Smith, C.J., Jeffs, A.G., Radford, C.A. (2013). Environmental influences on the larval recruitment dynamics of snapper, *Chrysophrys auratus* (Sparidae). *Marine and Freshwater Research*, 64(8): 726–740.
- Sinclair, F. (1999). *Kauwaeranga in Context*. Wellington: Victoria University. Retrieved from <http://www.kennett.co.nz/law/indigenous/1999/34.html>.
- Stanton, B., Sutton, P. (2003). Velocity measurements in the East Auckland Current northeast of North Cape, New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 37(1): 195–204.
- Statistics New Zealand. (2014). Annual report of Statistics New Zealand for the year ended 30 June 2014. Wellington: Statistics New Zealand. 79.
- Stephens, S. (2003). Ecological sustainability assessment for Firth of Thames shellfish aquaculture: Task 1 – Hydrodynamic modelling. Auckland Regional Council Technical Publication No. 252 / Environment Waikato Technical report 05/05. 34.
- Stevenson, M.L., Field, K.D., Holton, A.L. (1987). Regional background discussion paper on areas to be investigated for

- proposed marine protected areas in the Central Fisheries Management Area. Wellington: Ministry of Agriculture and Fisheries. 98.
- Stewart, S. (2011). Evidence of age-class truncation in some exploited marine fish populations in New South Wales, Australia. *Fisheries Research*, 108 (1): 209–213.
- Stockin, K.A., Pierce, G.J., Binedell, V., Wiseman, N., Orams, M.B. (2008). Factors affecting the occurrence and demographics of common dolphins (*Delphinus sp.*) in the Hauraki Gulf, New Zealand. *Aquatic Mammals*, 34(2): 200–211.
- Sutton, P.J.H., Roemmich, D. (2001). Ocean temperature climate off north-east New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 35: 553–565.
- Swales, A., Hume, T. (1995). Sedimentation history and potential future impacts of production forestry on the Wharekawa Estuary, Coromandel Peninsula. Prepared for Carter Holt Harvey Forests Ltd. NIWA report no. CHH004.
- Swales, A., Hume, T.M., Oldman, J.W., Green, M.O. (1997). Holocene sedimentation and recent human impacts in a drowned valley estuary. In Lumsden, J. (Ed.), *Proceedings of the 13th Australian Coastal and Ocean Engineering Conference*. Centre for Advanced Engineering, University of Canterbury, Christchurch, New Zealand. 895–900
- Swales, A., Williamson, R.B., Van Dam, L.F., Stroud, M., McGlone, M.S. (2002). Reconstruction of urban stormwater contamination of an estuary using catchment history and sediment profile dating. *Estuaries*, 25(1): 43–56.
- Talman, S., Norkko, A., Thrush, S., Hewitt, J. (2004). Habitat structure and the survival of juvenile scallops *Pecten novaezelandiae*: comparing predation in habitats with varying complexity. *Marine Ecology Progress Series*, 269: 197–207.
- Taylor, R.B., Morrison, M.A. (2008). Soft-sediment habitats and fauna of Omaha Bay, north-eastern New Zealand. *Journal of the Royal Society of New Zealand*, 38: 187–214.
- Taylor, R.B., Morrison, M.A., Shears, N. (2011). Establishing baselines for recovery in a marine reserve (Poor Knights Islands, New Zealand) using recollections of long-term divers. *Biological Conservation*, 144(12): 3038–3046.
- Tezanos-Pinto, G. (2009). Population Structure, Abundance and Reproductive Parameters of Bottlenose Dolphins (*Tursiops truncatus*) in the Bay of Islands (Northland, New Zealand). PhD. thesis, School of Biological Sciences, The University of Auckland. 243.
- Tezanos-Pinto, G., Constantine, R., Brooks, L., Jackson, J.A., Mourão, F., Wells, S., Baker, C.S. (2013). Decline in local abundance of bottlenose dolphins (*Tursiops truncatus*) in the Bay of Islands, New Zealand. *Marine Mammal Science*, 29(4): 390–410.
- Thomas, H.L., Shears, N.T. (2013). Marine Protected Area Networks: A comparison of approaches. The Royal Forest and Bird Protection Society of New Zealand, Wellington, New Zealand. Retrieved from http://202.160.118.26/files/file/MPA%20network%20FINAL_July29.pdf
- Thrush, S., Hewitt, J., Cummings, V., Dayton, P., Cryer, M., Turner, S., Funnell, G., Budd, R., Milburn, C., Wilkinson, M. (1998). Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecological Applications*, 8(3): 866–879.
- Thrush, S.F., Dayton, P.K. (2002). Disturbance to marine benthic habitats by trawling and dredging: Implications for marine biodiversity. *Annual Review of Ecology and Systematics*, 33: 449–473.
- Thrush, S.F., Hewitt, J.E., Cummings, V.J., Dayton, P.K. (1995). The impact of habitat disturbance by scallop dredging on marine benthic communities: what can be predicted from the results of experiments? *Marine Ecology Progress Series*, 129(1–3): 141–150.
- Thrush, S.F., Hewitt, J.E., Cummings, V.J., Dayton, P.K., Cryer, M., Turner, S.J., Funnell, G., Budd, R., Milburn, C., Wilkinson,

- M.R. (1998). Disturbance of the marine benthic habitat by commercial fishing: impacts at the scale of the fishery. *Ecological Applications*, 8: 866–879.
- Tian, R.C., Chen, C., Stokesbury, K.D.E., Rothschild, B., Cowles, G.W., Xu, Q., Hu, S., Harris, B.P., & Marino II, M.C. (2009a). Modelling the connectivity between sea scallop populations in the Middle Atlantic Bight and over Georges Bank. *Marine Ecology Progress Series*, 380: 147–160.
- Tian, R.C., Chen, C., Stokesbury, K.D.E., Rothschild, B.J., Cowles, G.W., Xu, Q., Hu, S., Harris, B.P., & Marino II, M.C. (2009b). Dispersal and settlement of sea scallop larvae spawned in the fishery closed areas on Georges Bank. *ICES Journal of Marine Science*, 66(10): 2155–2164.
- Tillin, H.M., Hiddink, J.G., Jennings, S., Kaiser, M.J. (2006). Chronic bottom trawling alters the functional composition of benthic invertebrate communities on a sea-basin scale. *Marine Ecology-Progress Series*, 318: 31–45.
- Torres, L.G. (2013). Evidence for an unrecognised blue whale foraging ground in New Zealand. *New Zealand Journal of Marine and Freshwater Research*, 47(2): 235–248.
- Townsend, A.J., de Lange, P.J., Duffy, C.A.J., Miskelly, C.M., Molloy, J., Norton, D.A. (2008). New Zealand threat classification system manual. Department of Conservation, Wellington. 35.
- Tuck, I.D., Hall, S.J., Robertson, M.R., Armstrong, E., Basford, D.J. (1998). Effects of physical trawling disturbance in a previously unfished sheltered Scottish sea loch. *Marine Ecology Progress Series*, 162: 227–242.
- Tuck, I.D., Hewitt, J.E. (2013). Monitoring change in benthic communities in Spirits Bay. *New Zealand Aquatic Environment and Biodiversity Report 111*. Wellington: Ministry for Primary Industries. 52.
- Tuck, I.D., Hewitt, J.E., Handley, S.J., Lundquist, C.J. (in press). Assessing the effects of fishing on soft sediment habitat, fauna and process. *New Zealand Aquatic Environment and Biodiversity Report*.
- Tupper, M., Boutilier, R.G. (1995). Effects of habitat on settlement, growth and post settlement survival of Atlantic cod (*Gadus morhua*). *Canadian Journal of Fisheries and Aquatic Sciences*, 52: 1834–41.
- Turner, S., Schwarz, A. (2006). Management and conservation of seagrass in New Zealand: an introduction. *Science for Conservation*, 264: 90.
- Turner, S.J. (1995). Restoring seagrass systems in New Zealand. *Water & Atmosphere*, 3: 9–11.
- Turner, S.J., Hewitt, J.E., Wilkinson, M.R., Morrisey, D.J., Thrush, S.F., Cummings, V.J., Funnell, G. (1999). Seagrass patches and landscapes: the influence of wind-wave dynamics and hierarchical arrangements of spatial structure on macrofaunal seagrass communities. *Estuaries*, 22(4): 1016–1032.
- Turner, S.J., Schwarz, A. (2004). Information for the management and conservation of seagrass in New Zealand. Prepared by NIWA for Department of Conservation. 48.
- van Houte-Howes, K.S.S., Turner, S.J., Pilditch, C.A. (2004). Spatial differences in macroinvertebrate communities in intertidal seagrass habitats and unvegetated sediment in three New Zealand estuaries. *Estuaries and Coasts*, 27(6): 945–957.
- Vant, B. (2011). Water Quality of the Hauraki Rivers and Southern Firth of Thames, 2000–09. Waikato Regional Council Technical Report, TR 2013/20.
- Visser, I. N. (2000). Orca (*Orcinus orca*) in New Zealand waters. Auckland: University of Auckland, School of Environmental and Marine Science. 194.
- Waikato Regional Council. (2006). Waikato Region Economy-Environment Futures Report, 2006. Technical Report 2006/51. WRC, Hamilton.
- Waikato Regional Council. (2012). Potential for effects from onsite wastewater in the Waikato region, with particular

- focus on development south and east of Hamilton: WRC Technical Report 2012/09. WRC, Hamilton. Retrieved from <http://www.waikatoregion.govt.nz/PageFiles/23921/TR201209.pdf>
- Waikato Regional Council. (2015). Harbour and Catchment Management Plan – Whangapoua, WRC Technical Report 2015/03. WRC, Hamilton. Retrieved from <http://www.waikatoregion.govt.nz/PageFiles/40291/TR201503.pdf>
- Waikato Regional Council. (2016). Earthworks - erosion and sediment control. [webpage] Retrieved from <http://www.waikatoregion.govt.nz/earthworks/>
- Waikato Regional Council. (n.d.). Create your own wetland plan. [webpage] Retrieved from <http://www.waikatoregion.govt.nz/Environment/Natural-resources/Water/Freshwater-wetlands/Restoring-a-wetland/Create-your-own-wetland-plan/>
- Waitangi Tribunal. (1995). Te Whanganui-a-Orotu Report. Wellington.
- Waitangi Tribunal. (1998). Te Ika Whenua Rivers Report. Wellington.
- Watling, L., Hoofd, M., Boulanger, M., Ferguson, N., Nouvian, C. (2014). The bottom line on trawling: How much more science do we need? Presented at the ICES Symposium on the effects of fishing on benthic fauna, habitat and ecosystem function, Tromsø, Norway.
- Watling, L., Norse, E.A. (1998). Disturbance of the seabed by mobile fishing gear: a comparison to forest clearcutting. *Conservation Biology*, 12(6): 1180–1197.
- Waycott, M., Duarte, C.M., Carruthers, T.J., Orth, R.J., Dennison, W.C., Olyarnik, S., Williams, S.L. (2009). Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proceedings of the National Academy of Sciences*, 106(30): 12377–12381.
- White, J.W., Botsford, L.W., Moffitt, E.A., Fischer, D.T. (2010). Decision analysis for designing marine protected areas for multiple species with uncertain fishery status. *Ecological Applications*, 20(6): 523–1541.
- Widdows, J., Pope, N.D., Brinsley, M.D., Asmus, H., Asmus, R.M. (2008). Effects of seagrass beds (*Zostera noltii* and *Z. marina*) on near-bed hydrodynamics and sediment re suspension. *Marine Ecology Progress Series*, 358: 125–136.
- Willis, T.J. (2013). Scientific and biodiversity values of marine reserves: a review. DOC Research and Development Series 340. Department of Conservation, Wellington. 70.
- Wiseman, N., Parsons, S., Stockin, K.A., Baker, C.S. (2011). Seasonal occurrence and distribution of Bryde’s whales in the Hauraki Gulf, New Zealand. *Marine Mammal Science*, 27(4): E253–E267.
- Woods, C., Schiel, D.R. (1997). Use of seagrass *Zostera novazelandica* (Setchell, 1933) as habitat and food by the crab *Macrophthalmus hirtipes* (Heller, 1862) (Brachyura: Ocypodidae) on rocky intertidal platforms in southern New Zealand. *Journal of Experimental Marine Biology and Ecology*, 214(1): 49–65.
- Worm, B., Barbier, E.B., Beaumont, N., Duffy, E., Folke, C., Halpern, B.S., Jackson, J.B.C., Lotze, H.K., Micheli, F., Palumbi, S.R., Sala, E., Selkoe, K.A., Stachowicz, J.J., Watson, R. (2006). Impacts of biodiversity loss on ocean ecosystem services. *Science*, 314: 787–90.
- Wyatt, S. (2011). Economic Impact of Coromandel Aquaculture. Sapere Research Group report prepared for the Hauraki-Coromandel Development Group. Retrieved from http://www.srgexpert.com/Coromandel%20EIS_Final_9%20Sept%202011.pdf
- Zaeschmar, J.R., Dwyer, S.L., Stockin, K.A. (2013). Rare observations of false killer whales (*Pseudorca crassidens*) cooperatively feeding with common bottlenose dolphins (*Tursiops truncatus*) in the Hauraki Gulf, New Zealand. *Marine Mammal Science*, 29(3): 555–562.
- Zeldis, J., Gall, M., Greig, M., Pinkerton, M., Richardson, K. (2001). La Niña shuts down upwelling in north eastern New Zealand. In: Changes in the marine environment. Proceedings of the Annual Conference of the Australian Marine

Sciences Association run in association with the New Zealand Marine Sciences Society, Townsville, Australia (3–6 July 2001). 39.

Zeldis, J., Oldman, J., Ballara, S., Richards, L. (2005). Physical fluxes, pelagic ecosystem structure, and larval fish survival in Hauraki Gulf, New Zealand. *Canadian Journal of Fisheries and Aquatic Sciences*, 62: 593–610.

Zeldis, J.R. (2004). New and remineralised nutrient supply and ecosystem metabolism on the northeastern New Zealand continental shelf. *Continental Shelf Research*, 24: 563–581.

Zeldis, J.R., Francis, R.I.C.C. (1998). A daily egg production method estimate of snapper biomass in the Hauraki Gulf, New Zealand. *ICES Journal of Marine Science*, 55(3): 522–534.

Zeldis, J.R., Walters, R.A., Greig, M.J.N., Image, K. (2004). Circulation over the north-eastern New Zealand continental slope, shelf and adjacent Hauraki Gulf, from spring to summer. *Continental Shelf Research*, 24: 543–561.

